Hydrobiologia

The International Journal of Aquatic Sciences

Volume 520, 1 June 2004

Nutrient uptake rates by the alien alga *Undaria pinnatifida* (Phaeophyta) (Nuevo Gulf, Patagonia, Argentina) when exposed to diluted sewage effluent
A.I. Torres, M.N. Gil, J.L. Esteves 1–6

Paleolimnological evidence of change in a shallow, hypereutrophic lake: Upper Klamath Lake, Oregon, USA
J.M. Eilers, J. Kann, J. Cornett, K. Moser, A.St. Amand 7–18

The influence of flow impoundment and river regulation on the distribution of riverine macroinvertabrates at Mammoth Cave National Park, Kentucky, U.S.A.
S.A. Grubbs, J.M. Taylor 19–28

Armatalona gen. n. – a new genus of subfamily Aloninae (Anomopoda, Chydoridae), separated from genus *Alona* Baird, 1840
A.Y. Sinev 29–47

Spatial and temporal variability in intertidal assemblages dominated by the mussel *Brachidontes rodriguezii* (d'Orbigny, 1846)
M.L. Adami, A. Tablado, J.L. Gappa 49–59

Feeding habits and trophic relationships of fishes utilizing an impacted coastal habitat, Singapore
S. Hajisamae, L.M. Chou, S. Ibrahim 61–71

Evaluation of and insights from ALFISH: a spatially explicit, landscape-level simulation of fish populations in the Everglades
H. Gaff, J. Chick, J. Trexler, D. DeAngelis, L. Gross, R. Salinas 73–87

Phytoplankton size structure stability in a meso-eutrophic subtropical lake
Y. Kamenir, Z. Dubinsky, T. Zohary 89–104

Regional distribution of diatom assemblages in the headwater streams of Luxembourg
R. Rimet, L. Ector, H.M. Cauchie, L. Hoffmann 105–117

Rapid assessment indicator of wetland integrity as an unintended predictor of avian diversity
M.A. Stapanian, T.A. Waite, G. Krzys, J.J. Mack, M. Micacchion 119–126

Diversity and distribution of Tubificidae, Naididae, and Lumbriculidae (Annalida: Oligochaeta) in the Netherlands: an evaluation of twenty years of monitoring data
R.C. Nijboer, M.J. Wetzel, P.F.M. Verdonschot 127–141

Distribution of carbon between sediment and water in macrophyte dominated lowland streams
R.J. Wilcock, G.F. Croker 143–152

Density of red alder (*Alnus rubra*) in headwaters influences invertebrate and detritus subsidies to downstream fish habitats in Alaska
M.S. Wipfli, J. Musslewhite 153–163

The consortium of the sponge *Ephydatia fluviatilis* (L.) living on the common reed *Phragmites australis* in Lake Piediluco (central Italy)
E. Gaino, T. Lancioni, G. La Porta, B. Todini 165–178

Use of morphological characters for species separation within the genus *Artemia* (Crustacea, Branchiopoda)
G. Mura, B. Brecciaroli 179–188

Two new species of *Paraprionospio* (Polychaeta: Spionidae) from the Grand Caribbean region and comments of the genus status
V.H. Delgado-Blas 189–198

Short research note

Effects of evaluation, wave exposure, and year on the proportion of gametophytes and tetrasporophytes in *Mazzaella parksii* (Rhodophyta, Gigartinaceae) populations
R. Scrosati, B. Mudge 199–205

Electronic journals at
KluwerOnline
WWW.KLUWERONLINE.NL
Contact your librarian for more information